

Resonance Broadening Induced Nonlinear Saturation of Kinetic Alfvén Turbulence in the Interplanetary Plasma

M.V. Medvedev ^{*} [†] and P.H. Diamond [‡]

Physics Department, University of California at San Diego, La Jolla, California 92093-0319.

(published: Phys. Letters A **219** (5-6), August 1996, 293-298)

The saturation of ion cyclotron Alfvén turbulence excited by beam particles is investigated using resonance broadening theory. The stochastic scattering which decorrelates particles, includes both random acceleration by electric fields and a turbulent magnetic mirroring effect. Turbulent mirroring is shown to yield non-Gaussian corrections to the orbits even if the random electric and magnetic fields are Gaussian. The predicted steady-state turbulence level exhibits a peaked angular distribution, with a maximum near $\Theta \simeq 60^\circ$.

PACS numbers: 96.50Ek, 52.35Qz, 52.35Ra, 52.40Mj

Keywords: resonance broadening, cyclotron instability

It is well known that the interaction of the solar wind plasma with planetary magnetospheres and cometary plasmas results in a high level of wave activity [1–3]. Gyration solar wind ions are a possible agent for exciting high-frequency instabilities. It has been suggested [4,5] that these magnetic fluctuations result from ion cyclotron instability, which occurs when the solar wind plasma interacts with a beam of cometary ions. A significant effort has been made to understand the turbulence generated by comet-solar wind interactions using quasilinear theory [4–6] as well as by simulation studies [7–12]. Ions are implanted in the solar wind plasma by melting and photoionization, thus forming an unstable ‘bump-on-tail’ like (i.e. shifted Maxwellian) distribution in the solar wind frame [13,14]. In the far-upstream vicinity of a comet, ions form a ring distribution. When the phase velocity of a circularly polarized wave coincides with the parallel velocity of particles, i.e. there is an ion cyclotron resonance $\omega - k_{\parallel}v_{\parallel} = \Omega_i$ between the particles and the wave, they resonantly pump the wave. Using quasilinear theory, saturation is predicted to occur via plateau formation in the distribution function. This mechanism is rather slow, with a saturation rate of the inverse plateau formation time (i.e. a velocity diffusion time). Saturation via resonance broadening is much faster, with a time scale comparable to the instability growth rate (i.e. since the resonance width is proportional to the fluctuation amplitude). Thus, even if macroscopic modification of the distribution function (à la plateau formation) occurs, it should be calculated in the presence of a finite width, fluctuation-broadened resonance. Moreover, the quasilinear theory predicts high level of turbulence at $\Theta \simeq 0^\circ$, only (Θ is the angle between the wave propagation direction and the ambient magnetic field), whereas observational data (for example, from the comet P/Grigg-Skjellerup [15,16]) indicates an average value of Θ between $50^\circ - 60^\circ$ at large distances from the comet ($R > 400,000$ km. for P/GS) and $75^\circ - 90^\circ$ closer to the comet. The saturation level prediction given here, which is based on resonance broadening theory, yields a peaked Θ -profile with the maximum value at $\Theta_m \simeq 65^\circ$ for P/GS parameters. Note that near-normal angles of propagation can also result from excitation of left hand polarized waves by normal Doppler resonance [17] in the case when $v_{\parallel} \cos \Theta < v_A$. For Jupiter we predict $\Theta_m^{th} \simeq 60^\circ$ and observations from the Ulysses spacecraft give $\Theta_m^{exp} \simeq 55^\circ - 60^\circ$ [24].

We develop a nonlinear kinetic model, so it is natural to anzatz that the high-frequency turbulent background is composed of a large number of kinetic shear Alfvén waves which are excited in plasma by the ion cyclotron instability. The ion-cyclotron resonance becomes $\omega - k_{\parallel}v_{\parallel} - \Omega_i \sim k_{\parallel}(v_A - v_{beam}) - \Omega_i$. We keep $(k_{\perp}\rho_i)^2$ terms, however, since they can be significant at large angles $k_{\parallel}/k_{\perp} \ll 1$. An energetic particle immersed in a such turbulent background is subject to stochastic scattering which includes both random acceleration by electric fields ($\tilde{\mathbf{E}}$, $\mathbf{v}_{\parallel} \times \tilde{\mathbf{B}}_{\perp}$) as well as a stochastic analogue of magnetic mirroring ($\mathbf{v}_{\perp} \times \tilde{\mathbf{B}}_{\perp}$). Random electric fields result in the diffusion of guiding centers in the perpendicular plane. However, the stochastic mirroring effect produces parallel diffusion in velocity space, giving rise to superdiffusion and rapid decorrelation (i.e. $\langle x^2 \rangle \sim \tau^3$) of particle orbit guiding centers along the ambient magnetic field. Since turbulent mirroring is an intrinsically nonlinear effect, it results in non-Gaussian behavior orbit fluctuation statistics even if the turbulent electric and magnetic fields obey Gaussian statistics. The random Doppler

^{*}E-mail address: mmedvedev@ucsd.edu

[†]Also: Institute for Nuclear Fusion, Russian Research Center “Kurchatov Institute”, Moscow 123182, RUSSIA.

[‡]Also: General Atomics, San Diego, California 92122.

shifts caused by this scattering decorrelate the particles from resonance. To calculate this stochastic decorrelation, we employ a modified stochastic orbit theory [18,19] which was developed for both low-frequency ($\omega \ll \Omega_i$) instabilities [20,21] and high-frequency ($\omega \simeq \Omega_i$) instabilities [22]. As is shown below, the theory yields results which are in very good agreement with the experimental data, in regards amplitude and $k_\perp/k_\parallel$ dependence.

Starting from the Vlasov equation, we separate the distribution function into the averaged (over fast time and space scales) part $\langle f(\mathbf{x}, \mathbf{v}, t) \rangle$, the test wave response $f_{\mathbf{k}}(\mathbf{x}, \mathbf{v}, t)$, and background response. We introduce electric and magnetic fields of the test wave in terms of scalar and vector potentials as usual, noting $\tilde{B}_\parallel = 0 \rightarrow \mathbf{A}_\perp = 0$. This follows from the fact that kinetic shear Alfvén waves are incompressible. Formally applying the procedure developed in [22], we now arrive at the lowest order solution of the Vlasov equation

$$f_{\mathbf{k}} = \frac{q}{m} \phi_{\mathbf{k}} \left\{ \mathcal{K}_\perp \frac{1}{v_\perp} \frac{\partial}{\partial v_\perp} + i \exp(-i\mathbf{k} \cdot \mathbf{x}) \sum_{n=-\infty}^{\infty} J_n^2 \left(\frac{k_\perp v_\perp}{\Omega_i} \right) R(\hat{\omega}) \left[(\omega - k_\parallel v_\parallel) \mathcal{K}_\perp \frac{1}{v_\perp} \frac{\partial}{\partial v_\perp} + \mathcal{K}_\parallel k_\parallel \frac{\partial}{\partial v_\parallel} \right] F_M \right\}, \quad (1)$$

where $\mathcal{K}_{\parallel, \perp}$ are the factors: $\mathcal{K}_\parallel = [1 - (1 - k_\perp v_\perp / \omega)(1 + k_\perp^2 \rho_s^2)]$, $\mathcal{K}_\perp = [1 - k_\parallel v_\parallel / \omega(1 + k_\perp^2 \rho_s^2)]$, and $\rho_s = c_s / \Omega_i$, c_s is the sound speed, $\Omega_i = eB_0 / m_i c$ is the ion cyclotron frequency. Note that in this formula the resonance function $R(\hat{\omega})$ is defined by the Fourier transform

$$R(\hat{\omega}) = \int_0^\infty d\tau \exp(i\hat{\omega}\tau) \langle \exp[i\mathbf{k} \cdot \delta\mathbf{x}(t - \tau)] \rangle \quad (2)$$

of the perturbed orbit propagator and asymptotes to the usual resonant denominator of the quasilinear theory when fluctuating turbulent forces vanish. Here $\delta\mathbf{x}$ is the deviation of a particle orbit from the unperturbed one, i.e. $\delta\mathbf{x}(t - \tau) = \mathbf{x}'(t - \tau) - \mathbf{x}(t - \tau)$ and $\hat{\omega} = \omega - k_\parallel v_\parallel - n\Omega_i$. Using Eq. (1), we may find (from the dispersion relation) the linear growth rate of ion cyclotron instability

$$\gamma_0 = -\frac{n_b}{n_0} \frac{T_i}{T_b} \frac{(\omega - \Omega_i)^2}{|k_\parallel| v_{T_b}} \left(\frac{\pi}{2} \right)^{1/2} \exp \left\{ -\frac{(\omega - \Omega_i)^2}{2k_\parallel^2 v_{T_b}^2} \right\} \left[1 - \frac{v_D}{v_A} (1 + k_\perp^2 \rho_s^2)^{1/2} - \frac{k_\perp^2 \rho_s^2}{\Omega_i} (\omega - k_\parallel v_D) \right], \quad (3)$$

where n_0 and T_i are the density and temperature of a bulk plasma, n_b and T_b are those of the beam, $\mathbf{v}_D$ is the longitudinal beam drift velocity such that $((\omega - k_\parallel v_D - \Omega_i) / k_\parallel v_{T_b})^2 \ll 1$, and v_{T_b} is the thermal velocity of the beam particles.

To explore the finite amplitude regime, note that the resonance function R is expressed via the perturbation of the orbit function of a test particle, Eq. (2). Expanding $\langle \exp(i\mathbf{k} \cdot \delta\mathbf{x}) \rangle$ in a Taylor series, with the conditions $\Omega_i \tau \gg 1$ and $\tau \ll \tau_{kinetic}$ ($\tau_{kinetic}$ is the characteristic time of the quasilinear diffusion), we have

$$\begin{aligned} \langle \exp(i\mathbf{k} \cdot \delta\mathbf{x}) \rangle &= \exp \left\{ \langle i\mathbf{k} \cdot \delta\mathbf{x} \rangle - \frac{1}{2!} \langle (\mathbf{k} \cdot \Delta\delta\mathbf{x})^2 \rangle + \dots \right\} \\ &= \exp \{ i\delta\omega\tau - k_\perp^2 D_\perp \tau - k_\parallel^2 D_{v_\parallel} \tau^3 + \dots \}, \end{aligned} \quad (4)$$

where $\Delta\delta\mathbf{x} = \delta\mathbf{x} - \langle \delta\mathbf{x} \rangle$, $\Delta\omega_\perp = k_\perp^2 D_\perp$, $D_\perp$ is the perpendicular diffusion coefficient and $D_{v_\parallel}$ is the diffusion coefficient of the longitudinal velocity in velocity space. The cubic in τ term gives rise to a Γ -function in the real-space representation of the resonance function. For the intermediate time-scale $\Omega_i^{-1} \ll \tau \ll \tau_{kinetic}$ as above, one can simplify the result by assuming $\tau^3 k_\parallel^2 D_{v_\parallel} \rightarrow \tau (k_\parallel^2 D_{v_\parallel})^{1/3}$. The resonance frequency shift $\delta\omega$ is proportional to spatial derivatives of the guiding center distribution, and vanishes for a Maxwellian [22]. Then, using the approximations mentioned above, the resonance function becomes

$$R(\hat{\omega}) = \frac{i}{\hat{\omega} + i\Delta\omega_\perp + i\Delta\omega_\parallel}, \quad (5)$$

where $\Delta\omega_\perp = k_\perp^2 D_\perp$ and $\Delta\omega_\parallel = (k_\parallel^2 D_{v_\parallel})^{1/3}$.

The system of equations describing the perturbed motion of a test particle cannot be solved exactly because of existence of the nonlinear coupling term $\mathbf{v} \times \mathbf{B}$. We use a perturbation method, assuming small perturbations. Up to the second order in perturbation, we have

$$\begin{aligned} \mathbf{x}(t + \tau) &= \mathbf{x}(t) + \hat{\mathbf{e}}_\perp \left[\frac{v_\perp}{2} \exp[i\Delta\psi] \frac{\exp(i\Omega_i\tau) - 1}{i\Omega_i} + c.c. \right] + \hat{\mathbf{e}}_\parallel v_\parallel \tau \\ &+ \hat{\mathbf{e}}_\perp \left[\frac{1}{2} \int_0^\tau ds \frac{q}{m} \left(\tilde{\mathcal{E}}^+(t + s) + \tilde{\mathcal{F}}^+(t + s) \right) \exp(-i\psi) \frac{\exp[i\Omega_i(\tau - s)] - 1}{i\Omega_i} + c.c. \right] \\ &+ \hat{\mathbf{e}}_\parallel \int_0^\tau ds (\tau - s) \frac{q}{m} \left(\tilde{\mathcal{E}}_\parallel(t + s) + \tilde{\mathcal{F}}_\parallel(t + s) \right), \end{aligned} \quad (6)$$

where $\Delta\psi$ is the angle between $\mathbf{k}_\perp$ and $\mathbf{v}_\perp$ and

$$\begin{aligned}\tilde{\mathcal{E}}_\perp^\pm(t) &= \tilde{E}_\perp^\pm(t) \mp i \frac{v_{\parallel 0}}{c} \tilde{B}_\perp^\pm(t), \quad \tilde{\mathcal{E}}_\parallel(t) = \tilde{E}_\parallel(t) + \frac{i}{2c} (v_{\perp 0}^+ \tilde{B}_\perp^-(t) - v_{\perp 0}^- \tilde{B}_\perp^+(t)), \\ \tilde{\mathcal{F}}_\perp^\pm(t) &= \mp i \frac{\tilde{v}_\parallel}{c} \tilde{B}_\perp^\pm(t), \quad \tilde{\mathcal{F}}_\parallel(t) = \frac{i}{2c} (\tilde{v}_\perp^+ \tilde{B}_\perp^-(t) - \tilde{v}_\perp^- \tilde{B}_\perp^+(t)), \\ \tilde{v}_\parallel(t) &= \frac{q}{m} \int_0^t \tilde{E}_\parallel(t') dt', \quad \tilde{v}_\perp^\pm(t) = e^{-i\Omega_i t} \int_0^t e^{i\Omega_i t'} \frac{q}{m} [\tilde{E}_\perp^\pm(t') \mp i \frac{v_{\parallel 0}}{c} \tilde{B}_\perp^\pm(t')] dt'.\end{aligned}\tag{7}$$

Here $\hat{\mathbf{e}}_\perp$ is the unit vector in the plane perpendicular to the ambient magnetic field parallel to $\mathbf{v}_\perp$, $\hat{\mathbf{e}}_\parallel$ is the unit vector collinear to the ambient magnetic field, $v_\perp^\pm = v_x \pm i v_y$ and $v_{\perp 0}^\pm, v_{\parallel 0}$ are the zeroth order velocities. From here we see that even if the $\tilde{\mathcal{E}} \propto \phi_{\mathbf{k}}$ field is Gaussian, $\tilde{\mathcal{F}}$ is not always Gaussian, but rather given by a distribution proportional to a Gaussian multiplied by an error function. This means that the third and other moments in the expansion (4) are non-zero. The error function is an increasing function of its argument. Thus, corrections appear on the tail of the Gaussian distribution, i.e. at sufficiently large times such that $\tau \sim \tau_{kinetic}$. This effect implies temporal intermittency for strongly mirrored particles. On intermediate time-scales $\Omega_i^{-1} \ll \tau \ll \tau_{kinetic}$, this non-Gaussian correction is small even when $\delta B/B \sim 1$. To capture the simple physical effect of resonance broadening we can drop the non-Gaussian terms $\tilde{\mathcal{F}}$ as small perturbations. After straightforward calculation using Eq. (4,6) and omitting rapidly oscillating terms in τ , we obtain

$$\begin{aligned}\Delta\omega_\perp &= \frac{k_\perp^2 v_A^2}{B_0^2 (1 + k_\perp^2 \rho_s^2)} \sum_{\mathbf{k}'} \sum_{n=-\infty}^{\infty} |B_{\perp \mathbf{k}'}|^2 \left\{ 1 + \frac{v_\parallel}{v_A} (1 + k_\perp'^2 \rho_s^2)^{1/2} \right\} \\ &\quad \cdot \frac{1}{4} [J_{n-1}^2(s) + 2J_n^2(s) + J_{n+1}^2(s)] R(\omega - k_\parallel v_\parallel - n\Omega_i)\end{aligned}\tag{8}$$

and similarly

$$\begin{aligned}\Delta\omega_\parallel &= \left\{ \frac{\Omega_i^2 k_\parallel^4}{B_0^2 k_\perp^2} \frac{v_A^2}{(1 + k_\perp^2 \rho_s^2)} \sum_{\mathbf{k}'} \sum_{n=-\infty}^{\infty} |B_{\perp \mathbf{k}'}|^2 \left(J_n^2(s) \right. \right. \\ &\quad \left. \left. + \frac{1}{4} \frac{k_\perp'^2 v_\perp^2}{k_\parallel'^2 v_A^2} [J_{n-1}^2(s) - J_{n+1}^2(s)] R(\omega - k_\parallel v_\parallel - n\Omega_i) \right) \right\}^{1/3}.\end{aligned}\tag{9}$$

The sums in these equations run over the full range of the background turbulence spectrum $-\infty < k' < \infty$. Here ‘unprimed’ wave vectors refer to the test wave. Eqs. (8,9) describe the general case of resonance broadening due to the effects of scattering of a test particle by random fields. They can be simplified by assuming a narrow-band spectrum [22]. As it will be shown later from estimating the spectrum at saturation, the narrow spectrum approximation is valid for this case. Then $\sum_{\mathbf{k}} |B_{\perp \mathbf{k}}|^2 / B_0^2$ can be replaced by $(\delta B/B)^2$. As we consider the ion cyclotron resonance with ions we may set $n = 1$. $n \neq 1$ terms give only a small correction.

The broadening $\Delta\omega(\mathbf{v})$ should be averaged over the velocity distribution function [22]. Saturation occurs when the resonance broadening becomes comparable to the linear growth rate γ_0 , i.e. when the total nonlinear growth rate vanishes, $\gamma_{NL} = \gamma_0 - \overline{\Delta\omega} = 0$. This saturation mechanism does not rely upon quasilinear plateau formation and thus is applicable to instances where the unstable distribution is maintained externally, such as in the case of comet-solar wind interaction. For the ion cyclotron instability, we define the average broadening $\overline{\Delta\omega} = \Gamma_1^{-1}(s_i) \int d\mathbf{v} F_M(\mathbf{v}) J_1^2(k_\perp v_\perp / \Omega_i) \Delta\omega(\mathbf{v})$, where $s_i = k_\perp v_{Ti} / \Omega_i$ and $\Gamma_n(s_i) = \int d\mathbf{v} F_M(\mathbf{v}) J_n^2(k_\perp v_\perp / \Omega_i)$, $n = 0, 1, \dots$. Upon averaging of Eqs. (8,9) with $R(\tilde{\omega})$ written for the unperturbed case, we can calculate the nonlinear saturation level. We use a perturbation analysis to explore those regions where the turbulence level is (sufficiently) small, i.e. at angles of propagation close to 0° and 90° (see below). For the case of the perpendicular diffusion (i.e. $\gamma_0 - \Delta\omega_\perp = 0$), we obtain

$$\left(\frac{\delta B}{B} \right)_{NL\perp}^2 = \frac{(\omega - k_\parallel v_D - \Omega_i)^2 (1 + k_\perp^2 \rho_s^2)}{k_\perp^2 v_A^2 \left(1 + \frac{v_D}{v_A} (1 + k_\perp^2 \rho_s^2)^{1/2} \right) F_1(s_i)},\tag{10}$$

where $v_A^2 = B_0^2 / 4\pi\rho$ is the Alfvén speed, $F_1(s_i) = \Gamma_1^{-1}(s_i) \int 0.5 J_1^2(J_0^2 + 2J_1^2 + J_2^2) F_M(\mathbf{v}) d\mathbf{v}$. The saturation level is roughly proportional to $k_\perp^{-2} \sim \sin^{-2} \Theta$. Thus this case corresponds to the saturation at large angles. By contrast, if there is only parallel diffusion (i.e. $\gamma_0 - \Delta\omega_\parallel = 0$), we have




Fig. 1




Fig. 2

FIG. 1. The angular distribution of the saturated Alfvén turbulence fluctuation level.

FIG. 2. The angular distribution of the MHD activity level (in arbitrary units) near Jupiter obtained from Ulysses data.

$$\left(\frac{\delta B}{B}\right)_{NL\parallel}^2 = \frac{\gamma_0^2 (\omega - k_{\parallel} v_D - \Omega_i)^2 (1 + k_{\perp}^2 \rho_s^2) k_{\perp}^2}{\Omega_i^2 k_{\parallel}^4 v_A^2 \left[F_2(s_i) + 2 \frac{k_{\perp}^2 v_{T_i}^2}{k_{\parallel}^2 v_A^2} F_3(s_i) \right]}, \quad (11)$$

where $F_2(s_i) = \Gamma^{-1}(s_i) \int J_1^4 F_M(\mathbf{v}) d\mathbf{v}$, $F_3(s_i) = \Gamma^{-1}(s_i) \int 0.25 J_1^2 [J_0^2 - J_2^2] F_M(\mathbf{v}) v_{\perp} d\mathbf{v}$. The fluctuation level $\delta B/B$ in this case is proportional to $k_{\parallel}^{-2} \sim \cos^{-2} \Theta$. Thus parallel diffusion suppress turbulence at small angles. In the derivation of this equation we made the approximation $\overline{f^3} \approx \overline{f}^3$, but indeed they differ only by a factor close to unity. The $\Delta\omega_{\parallel}$ effect is dominant at small angles of wave propagation, whereas the $\Delta\omega_{\perp}$ effect is more important at near-perpendicular angles. For the case where $\Delta\omega = \Delta\omega_{\perp} + \Delta\omega_{\parallel}$ it is necessary to solve the third order algebraic equation of the type $\gamma_0 = c_1 \delta B^2 + c_2 \delta B^{2/3}$. Fig. 1 represent the solution of this equation for typical comet and solar wind parameters: $n = 10 \text{ cm}^{-3}$, $n_b = 1.7 \text{ cm}^{-3}$, $T_e = T_i = 10 \text{ eV}$, $T_b = 3 \text{ eV}$, $v_{T_i} = 3 \cdot 10^7 \text{ cm/s}$, $v_A = 0.2 v_{T_i}$, $v_D = 2 v_A$, $\Omega_i = 10 \text{ s}^{-1}$, $k = 10^{-5} \text{ cm}^{-1}$.

Fig. 1 shows the anisotropy of magnetic fluctuation level in propagation angle Θ ($\tan \Theta = k_{\perp}/k_{\parallel}$). One can see that at small angles ($\Theta \simeq 0^\circ$) with respect to the ambient magnetic field, very small background turbulence levels saturate the ion cyclotron instability. The same is true of the near-perpendicular case, $\Theta \sim 90^\circ$. There the turbulence is strongly self-suppressed. This is not the case for $\Theta \simeq 60^\circ$. Here, the amplitude of the background Alfvén waves must be large for resonance broadening to self-consistently suppress the ion cyclotron instability. The main energy content of the observable MHD activity, therefore, is associated with Alfvén waves propagating at large angles relative to the averaged magnetic field. The peak is strongly localized at angles near $\Theta_m \simeq 60^\circ$. The value Θ_m of the maximum turbulent fluctuation is remarkably insensitive to plasma parameters such as bulk plasma and ion beam temperatures, density and drift velocity of the beam, etc. For all reasonable parameters, Θ_m remains approximately 60 degrees. The magnitude of the maximum fluctuation level, on the contrary, depends sensitively on the beam and plasma parameters. Fig. 2 is a plot of experimental data of MHD activity near Jupiter, obtained by the spacecraft Ulysses [24]. Here the level of fluctuations (in arbitrary units), measured for different angles, is shown. It is apparent that the dominant waves are those propagating at angles in the interval $50^\circ < \Theta < 70^\circ$. There is remarkable agreement of the angular distribution obtained from experiments [15,16,24] with our very simple theory. Quasilinear theory predicts a different result [5,17], which is (approximately) $(m_i n_i / m_p n_p)^{2/3} (v_D / v_A)^{1/3}$. The Θ -dependence predicted by quasilinear theory is not in agreement with experimental data, although it gives a reasonable prediction of the magnitude, i.e. $\delta B/B \sim 1$.

We emphasize the fact that the peaked angular distribution is the result of both perpendicular and parallel diffusion. The perpendicular diffusion (the diffusion of guiding centers) is a result of the magnetic field line flutter and is responsible for the resonance broadening at larger angles, $\Theta \sim 90^\circ$. The parallel diffusion (the randomized step-size of a particle helical trajectory) is due to the random mirroring force. It is responsible for the instability suppression at small angles of wave propagation, i.e. for Θ near 0° . It is important to note that the parallel diffusion in velocity space results in a $\langle x^2 \rangle \sim \tau^3$ type decorrelation process in real space. However, for intermediate time-scales $\Omega_i^{-1} \ll \tau \ll \tau_{kinetic}$, it can be modelled by a standard diffusion process. The magnetic mirroring gives rise to

non-Gaussian dynamics of a particle immersed into turbulent background, even if this background is Gaussian. This describes temporal intermittancy in a system with strongly mirrored particles ($\delta B/B \sim 1$).

We are very grateful to S.K. Ride for discussions and for unpublished data from the Ulysses spacecraft. We also thank V.D. Sapiro, V.I. Shevchenko and B.T. Tsurutani for useful discussions. This research was supported by U.S. Department of Energy, Contract No. DE-FG03-88ER53275 and NASA Grant No. UT-A:NAGW-2418.

-
- [1] B.T. Tsurutani, E.J. Smith, *Geophys. Res. Lett.*, **13**, 263 (1986).
 - [2] W. Reidler, K. Schwingenschuh, Ye.G. Yeroshenko, V.A. Styashkin, C.T. Russell, *Nature*, **321**, 288 (1986).
 - [3] B.T. Tsurutani, P. Rodriguez, *J. Geophys. Res.*, **86**, 4319 (1981).
 - [4] R.Z. Sagdeev, V.D. Sapiro, V.I. Shevchenko, K. Szegö, *Geophys. Res. Lett.*, **13**, 85 (1986).
 - [5] A.A. Galeev, A.N. Polyudov, R.Z. Sagdeev, K. Szegö, V.D. Sapiro, V.I. Shevchenko, *Sov. Phys. JETP*, **65**, 1178 (1987).
 - [6] A.A. Galeev, *Exploration of Halley's Comet*, ESA SP-250, **1**, 3 (1986).
 - [7] K. Akimoto, D. Winske, *Phys. Rev. Lett.*, **64**, 753 (1990).
 - [8] K. Akimoto, D. Winske, S.P. Gary, M.F. Tomsen, *J. Geophys. Res.*, **98**, 1419 (1993).
 - [9] A.J. Coates, A.D. Johnstone, D.E. Huddleston, B. Wilken, *Geophys. Res. Lett.*, **20**, 483 (1993).
 - [10] S.P. Gary, R.H. Miller, D. Winske, *Geophys. Res. Lett.*, **18**, 1067 (1991).
 - [11] M. Hoshino, T. Tetasawa, *J. Geophys. Res.*, **90**, 57 (1985).
 - [12] T. Terasawa, *Phys. Commun.*, **49**, 193 (1988).
 - [13] C.F. Kennel, *J. Geophys. Res.*, **86**, 4325 (1981).
 - [14] E.W. Greenstadt, R.W. Fredericks, in *Solar Wind Plasma Physics III*, edited by C.F. Kennel, L.J. Lanzerotti, E.N. Parker, North-Holland, Amsterdam (1979).
 - [15] F.M. Neubauer, K.-H. Glassmeier, A.J. Coates, A.D. Johnstone, *J. Geophys. Res.*, **98**, 20937 (1993).
 - [16] F.M. Neubauer, H. Marshall, Pohl, K.-H. Glassmeier, G. Mushman, F. Mariani, M.H. Acuna, L.F. Burlaga, N.F. Ness, M.K. Wallis, H.V. Schmidt, E. Urgstrup, *Astron. Astrophys.*, **L5**, 268 (1993).
 - [17] V.I. Shevchenko, V.D. Shapiro, S.K. Ride, M. Baine, *J. Geophys. Res.*, **100**, 1735 (1995).
 - [18] T.H. Dupree, *Phys. Fluids*, **9**, 1733 (1966).
 - [19] R.E. Aamodt, *Phys. Fluids*, **10**, 1245 (1967).
 - [20] T.H. Dupree, *Phys. Fluids*, **10**, 1049 (1967).
 - [21] T.H. Dupree, *Phys. Fluids*, **11**, 2680 (1968).
 - [22] C.T. Dum, T.H. Dupree, *Phys. Fluids*, **13**, 2064 (1969).
 - [23] T.M. Antonsen, Jr., B. Lane, *Phys. Fluids*, **23**, 1205 (1980).
 - [24] S.K. Ride, *private communication* (1994).